

AD-A148 711

MULTIDIMENSIONAL INVERSE SCATTERING FOR FIRST ORDER  
SYSTEMS(U) CLARKSON UNIV POTSDAM NY INST FOR NONLINEAR  
STUDIES A I NACHMAN ET AL. JUL 84 AFOSR-TR-84-0918

1/1

UNCLASSIFIED

AFOSR-78-3674


F/G 12/1

NL

END

FORM

DTG



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A


2

I.F.N.S. #38

APOSR-TR. 84-0918

# INSTITUTE FOR NONLINEAR STUDIES

AD-A148 711



MULTIDIMENSIONAL INVERSE SCATTERING  
FOR FIRST ORDER SYSTEMS

by

Approved for public release;  
distribution unlimited.

Adrian I. Nachman\* and Mark J. Ablowitz  
Clarkson University

Clarkson College of Technology  
Potsdam, New York 13676

\*Permanent Address: Department of  
Mathematics, University of  
Rochester, Rochester, NY 14627

July 1984

84 10 23 30

DTIC FILE COPY

DTIC  
ELECTE  
DEC 06 1984  
S D  
E

AD-A148711

## REPORT DOCUMENTATION PAGE

1. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
3. SECURITY CLASSIFICATION / DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 84-0918</b>	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
NAME OF PERFORMING ORGANIZATION Clarkson University		6b. OFFICE SYMBOL (If applicable)	
ADDRESS (City, State, and ZIP Code) Dept of Mathematics & Computer Science Potsdam NY 13676		7b. ADDRESS (City, State, and ZIP Code) Directorate of Mathematical & Information Sciences, AFOSR, Bolling AFB DC 20332	
NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM	
ADDRESS (City, State, and ZIP Code) Bolling AFB DC 20332		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-78-3674	
10. SOURCE OF FUNDING NUMBERS		PROGRAM ELEMENT NO. 61102F	
		PROJECT NO. 2304	
TASK NO. A4		WORK UNIT ACCESSION NO.	
TITLE (Include Security Classification) MULTIDIMENSIONAL INVERSE SCATTERING FOR FIRST ORDER SYSTEMS			
12. PERSONAL AUTHOR(S) Adrian I. Nachman* and Mark J. Ablowitz			
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM TO	
14. DATE OF REPORT (Year, Month, Day) JUL 84		15. PAGE COUNT 14	
16. SUPPLEMENTARY NOTATION *Permanent address: Dept of Mathematics, Univ of Rochester, Rochester NY 14627.			
COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
9. ABSTRACT (Continue on reverse if necessary and identify by block number) A method for solving the inverse problem for a class of multidimensional first order systems is given. The analysis yields equations which the scattering data must satisfy; these equations are natural candidates for characterizing admissible scattering data. The results are used to solve the multidimensional N-wave resonant interaction equations.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
NAME OF RESPONSIBLE INDIVIDUAL Dr. Robert N. Buchal		22b. TELEPHONE (Include Area Code) (202) 767- 4939	
		22c. OFFICE SYMBOL NM	

# MULTIDIMENSIONAL INVERSE SCATTERING FOR FIRST ORDER SYSTEMS

by

Adrian I. Nachman\* and Mark J. Ablowitz  
Department of Mathematics and Computer Science  
Clarkson University, Potsdam, New York 13676

## ABSTRACT

A method for solving the inverse problem for a class of multidimensional first order systems is given. The analysis yields equations which the scattering data must satisfy; these equations are natural candidates for characterizing admissible scattering data. The results are used to solve the multidimensional N-wave resonant interaction equations.

### 1. Introduction

The inverse scattering problems for the hyperbolic and elliptic generalizations in the plane of the  $m \times m$  AKNS system have been successfully studied in [1] and applied to the linearization of several physically significant nonlinear evolution equations (N-wave interaction, Davey-Stewartson, etc.) in two spatial and one temporal dimensions.

We indicate here how the method used in our investigation of the  $n$ -dimensional Schrödinger equation [2] can be applied to the study of the inverse problem for the operator in  $\mathbb{R}^{n+1}$ :

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH  
NOTICE OF RESEARCH RESULTS  
This technical report is the result of research  
approved by the Air Force Office of Scientific Research  
Distribution Statement  
MATTHEW J. ABLOWITZ  
Chief, Technical Information Division

\*Permanent address: Department of Mathematics, University of Rochester, Rochester,  
New York 14627

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By	
Distribution /	
Availability Codes	
Avail and/or	
Dist	
A-1	

$$L_{\sigma} \doteq \frac{\partial}{\partial x_0} + \sigma \sum_{\ell=1}^n J_{\ell} \frac{\partial}{\partial x_{\ell}} - Q(x_0, x). \quad (1)$$

Here  $J_{\ell}$  are constant real diagonal  $m \times m$  matrices (we denote the diagonal entries of  $J_{\ell}$  by  $J_{\ell}^1, \dots, J_{\ell}^m$  and assume  $J_{\ell}^i \neq J_{\ell}^j \neq 0$  whenever  $i \neq j$ ); the matrix-valued off-diagonal potential  $Q = (Q^{ij})$  may depend on  $x_0$  as well as  $x = (x_1, \dots, x_n)$  and  $\sigma = \sigma_R + i\sigma_I$  is a complex parameter.

The operator (1) is associated with the nonlinear system:

$$\frac{\partial Q^{ij}}{\partial t} = \frac{1}{\sigma} a_{ij} \frac{\partial Q^{ij}}{\partial x_0} + \sum_{\ell} (a_{ij} J_{\ell}^i - B_{\ell}^i) \frac{\partial Q^{ij}}{\partial x_{\ell}} + \frac{1}{\sigma} \sum_{\ell} (a_{i\ell} - a_{\ell j}) Q^{i\ell} Q^{\ell j} \quad (2)$$

$$(\text{with } a_{ij} = \frac{B_{\ell}^j - B_{\ell}^i}{J_{\ell}^j - J_{\ell}^i}, \quad 1 \leq \ell \leq n, \text{ for some real } B_{\ell}^i \quad 1 \leq \ell \leq n, \quad 1 \leq i \leq m). \quad (3)$$

Even though no traditional scattering operator exists in the case  $\sigma_I \neq 0$ , the so-called  $\bar{\partial}$  method (see [2] and references given there) gives a satisfactory definition of scattering data for  $L_{\sigma}$ , along with a systematic inversion procedure, which we use to solve (2).

A solution of the inverse scattering problem for the hyperbolic case  $\sigma_I = 0$  is then obtained by a limiting argument; thus we can avoid a separate study of a Riemann-Hilbert boundary value problem (which in higher dimensions may also involve some geometric complications). The main advantage of this approach is that it yields (from the compatibility conditions associated with  $\bar{\partial}$  in several variables) equations which must be satisfied by the scattering data. In addition to their importance for the problem of characterizing admissible scattering data, these equations have several consequences: i) they provide a formula for reconstructing the potential from the scattering transform purely by quadratures (in the generic case when no three of the vectors  $J^i = (J_1^i, J_2^i, \dots, J_n^i)$ ,  $1 \leq i \leq m$ , are colinear); ii) they show how one can recover the scattering transform from (at least small) data given on certain  $(n+1)$ -dimensional surfaces ( $n+1$  being the number of variables in  $Q$ ); iii) they may indicate what other

(possibly non-local) evolution equations could be solvable with the IST developed here; iv) they constitute in themselves a new class of multidimensional nonlinear systems of integro-differential equations which are linearizable.

## 2. The Case $\sigma_I \neq 0$ .

We will denote by  $k = (k_1, \dots, k_n) = k_R + ik_I$  a point in  $\mathbb{C}^n$  and will often write  $f(k)$  instead of  $f(k_R, k_I)$  for an arbitrary function of  $k_R$  and  $k_I$ .

As a first step in the  $\bar{\partial}$  procedure we construct a family of solutions of  $L_\sigma \psi = 0$  of the form  $\psi = \mu(x_0, x, k) \exp[i\sigma \sum_{\ell=1}^n k_\ell (x_\ell - \sigma x_0 J_\ell)]$  with  $\mu$  bounded;  $\mu$  will then satisfy the equation

$$\frac{\partial \mu}{\partial x_0} + \sigma \sum_{\ell=1}^n J_\ell \frac{\partial \mu}{\partial x_\ell} + i\sigma \sum_{\ell=1}^n k_\ell [J_\ell, \mu] = Q\mu. \quad (4)$$

The generalized eigenfunctions  $\mu_\sigma = (\mu_\sigma^{ij})$  we will work with, are obtained by solving the integral equation  $\mu_\sigma = I + \tilde{G}_\sigma(Q\mu_\sigma)$ , i.e.

$$\mu_\sigma^{ij} = \delta_{ij} + \iint_{\mathbb{R}^{n+1}} G_\sigma^{ij}(x_0 - y_0, x - y, k) (Q(y_0, y) \mu_\sigma(y_0, y, k))^{ij} dy_0 dy, \quad (5)$$

where the Green's function is given by

$$G_\sigma^{ij}(x_0, x, k) = \frac{-i}{(2\pi)^{n+1}} \iint_{\mathbb{R}^{n+1}} \frac{e^{i(x_0 \xi_0 + x \cdot \xi)}}{\xi_0 + \sigma \sum_{\ell=1}^n [J_\ell^i \xi_\ell + k_\ell (J_\ell^i - J_\ell^j)]} d\xi_0 d\xi. \quad (6)$$

For brevity we will assume here that  $Q$  is such that this integral equation has a bounded solution  $\mu_\sigma$  for all  $k \in \mathbb{C}^n$ .

$G_\sigma$  can be computed explicitly by contour integration:

$$G_\sigma^{ij}(x_0, x, k) = \frac{\text{sign}(\sigma_I J_1^i)}{2\pi i (x_1 - \sigma J_1^i x_0)} e^{i\alpha_\sigma^{ij}(x_0, x, k)} \prod_{\ell=2}^n \delta(x_\ell - \frac{J_\ell^i}{J_1^i} x_1) \quad (7)$$

with

$$\alpha_\sigma^{ij}(x_0, x, k) = \sum_{\ell=1}^n \frac{J_\ell^i - J_\ell^j}{\sigma_I} (|\sigma|^2 x_0 k_{I_\ell} - \frac{x_\ell}{J_\ell^i} (\sigma k_\ell)_I). \quad (8)$$

The next step is to express  $\bar{\partial}\mu$  in terms of  $\mu$ . We start by writing

$\frac{\partial \tilde{G}}{\partial \bar{k}_p}$  and hence  $\frac{\partial \tilde{G}}{\partial \bar{k}_p} (Q\mu)$  as a superposition of exponentials:

$$\left(\frac{\partial \tilde{G}}{\partial \bar{k}_p} (Q\mu_\sigma)\right)^{ij} = \frac{\bar{\sigma}(J_p^i - J_p^j)}{2i|\sigma_I|(2\pi)^n} \int_{\mathbb{R}^n} \sigma\left(\sum_{\ell=1}^n J_\ell^i \lambda_\ell\right) e^{i\beta_\sigma^{ij}(x_0, x, k, \lambda)} T_\sigma^{ij}(k, \lambda) d\lambda. \quad (9)$$

$$\text{with } \beta_\sigma^{ij}(x_0, x, k, \lambda) = \alpha_\sigma^{ij}(x_0, x, k) + \sum_{\ell=1}^n (x_\ell - \sigma_R J_\ell^i x_0) \lambda_\ell \quad \text{and} \quad (10)$$

$$T_\sigma^{ij}(k, \lambda) = \iint_{\mathbb{R}^{n+1}} e^{-i\beta_\sigma^{ij}(y_0, y, k, \lambda)} (Q(y_0, y) \mu_\sigma(y_0, y, k))^{ij} dy_0 dy. \quad (11)$$

The calculation of  $\bar{\mu}$  is then based on the following crucial symmetry property of our Green's function:

$$e^{-i\beta_\sigma^{ij}(x_0, x, k, \lambda)} G_\sigma^{rj}(x_0, x, k) = G_\sigma^{ri}(x_0, x, \hat{k}_\sigma^{ij}(k, \lambda)) \text{ whenever } \sum_{\ell} J_\ell^i \lambda_\ell = 0; \quad (12)$$

here  $\hat{k}_\sigma^{ij}(k, \lambda)$  is the point in  $\mathbb{C}^n$  whose  $\ell^{\text{th}}$  component is

$$(\hat{k}_\sigma^{ij}(k, \lambda))_\ell = \frac{J_\ell^j - J_\ell^i}{\sigma_I J_\ell^i} (\sigma k_\ell)_I + k_\ell + \lambda_\ell. \quad (13)$$

Once (12) has been established it can be shown (assuming that (5) admits no homogeneous solutions) that

$$\begin{aligned} \frac{\partial \mu_\sigma}{\partial \bar{k}_p} (x_0, x, k) &= \sum_{i,j} \frac{\bar{\sigma}(J_p^i - J_p^j)}{2i|\sigma_I|(2\pi)^n} \int_{\mathbb{R}^n} \delta(\sum_{\ell} J_\ell^i \lambda_\ell) T_\sigma^{ij}(k, \lambda) e^{i\beta_\sigma^{ij}(x_0, x, k, \lambda)} \times \\ &\times \mu_\sigma(x_0, x, \hat{k}_\sigma^{ij}(k, \lambda)) E_{ij} d\lambda; \end{aligned} \quad (14)$$

(we have denoted by  $E_{ij}$  the  $m \times m$  matrix with entries  $E_{ij}^{rs} = \delta_{ir} \delta_{js}$ ). If we now fix all  $k_\ell$ ,  $\ell \neq p$ , and apply the (1-dimensional) inhomogeneous Cauchy integral formula



$$f(z) = \frac{1}{2\pi i} \int_{|z'|=R} \frac{f(z')}{z'-z} dz' + \frac{1}{2\pi i} \iint_{|z'| \leq R} \frac{\frac{\partial f}{\partial \bar{z}}(z')}{z'-z} dz' \wedge d\bar{z}' \quad (15)$$

to the  $k_p$  variable, we can convert (14)<sub>p</sub> to an integral equation: noting that  $\mu(x_0, x, k) \sim I$  when  $|k_p| \rightarrow \infty$  (and denoting  $k' = (k_1, \dots, k'_p, \dots, k_n)$ ) we have

$$\begin{aligned} \mu_\sigma(x_0, x, k) = I - \frac{i\bar{\sigma}}{|\sigma_I|(2\pi)^{n+1}} \sum_{i,j} (J_p^{i,j} - J_p^{j,i}) \iiint \frac{\delta(\Sigma J_\ell^{i,j} \lambda_\ell)}{k_p - k'_p} T_\sigma^{i,j}(k', \lambda) e^{i\beta_\sigma^{i,j}(x_0, x, k', \lambda)} \times \\ \times \mu_\sigma(x_0, x, \hat{k}^{i,j}(k', \lambda)) E_{ij} d\lambda dk'_R dk'_I. \end{aligned} \quad (16)_p$$

(More generally, one can use (15) with  $f(z) = \mu_\sigma(x_0, x, k+zv)$ ,  $z \in \mathbb{C}$ , with  $k$  fixed and with an arbitrary  $v \in \mathbb{C}^n$  which is not perpendicular to any of the vectors  $J^i - J^j$ ,  $i \neq j$ ). The matrix-valued function  $T_\sigma(k, \lambda)$  defined in (11) is our scattering data and (16) is the inverse scattering recipe for reconstructing  $\mu$  from  $T$ . Once  $\mu$  is found, the potential is easily recovered:

$$Q(x_0, x) = \frac{i\sigma}{\pi} [J_p, \iint \frac{\partial \mu_\sigma}{\partial \bar{k}_p}(x_0, x, k) dk_R dk_I]. \quad (17)$$

On the other hand, given an arbitrary  $T(k, \lambda)$ , to apply the above inversion procedure we would first need to know that the equations (14)<sub>p</sub>,  $p = 1, 2, \dots, n$ , are compatible; requiring that  $\frac{\partial^2 \mu}{\partial \bar{k}_r \partial \bar{k}_p} = \frac{\partial^2 \mu}{\partial \bar{k}_p \partial \bar{k}_r}$  yields the following characterization equations for  $T$ :

$$\begin{aligned} L_{pr}^{ij}[T_\sigma] \doteq (J_p^{i,j} - J_p^{j,i}) \frac{\partial T_\sigma^{ij}}{\partial \bar{k}_r} - (J_r^{i,j} - J_r^{j,i}) \frac{\partial T_\sigma^{ij}}{\partial \bar{k}_p} + \frac{i\bar{\sigma}}{2\sigma_I} (J_p^{i,j} - J_p^{j,i})(J_r^{i,j} - J_r^{j,i}) \left( \frac{1}{J_r^i} \frac{\partial T_\sigma^{ij}}{\partial \lambda_r} - \frac{1}{J_p^i} \frac{\partial T_\sigma^{ij}}{\partial \lambda_p} \right) = \\ = N_{pr}^{ij}[T_\sigma] \doteq \frac{i\bar{\sigma}}{2|\sigma_I|(2\pi)^n} \sum_{i'} [(J_p^{i'} - J_p^{j'}) (J_r^{i'} - J_r^{j'}) - (J_r^{i'} - J_r^{j'}) (J_p^{i'} - J_p^{j'})] \int \delta(\Sigma J_\ell^{i',j'} \lambda'_\ell) T_\sigma^{i',j'}(k, \lambda') \times \\ \times T_\sigma^{i',j'}(\hat{k}^{i',j'}(k, \lambda'), \lambda - \frac{J^{i'}}{J^i} \lambda') d\lambda'. \end{aligned} \quad (18)_{pr}^{ij}$$

For compatibility, (18)<sup>ij</sup> need only hold whenever  $\Sigma J_\ell^{i,j} \lambda_\ell = 0$ , however one may also

verify that  $T_\sigma$  when given by (11) satisfies (18) everywhere.

It turns out to be very useful to recast (18) in integral form. It is enough to keep only the equations (18)<sub>p1</sub>. We then look for a parametrization of the hyperplane  $\{(k, \lambda) \in \mathbb{C}^n \times \mathbb{R}^n : \sum J_\ell^i \lambda_\ell = 0\}$  by new variables  $(\chi, w_0, w) \in \mathbb{C}^{n-1} \times \mathbb{R} \times \mathbb{R}^n$  so that, in the new coordinates  $L_{p1}^{ij} = \frac{\partial}{\partial \chi_p}$ ,  $2 \leq p \leq n$  and

$\beta_\sigma^{ij}(\chi_0, \chi, k, \lambda) = \chi_0 w_0 + \chi \cdot w$ ; these requirements determine (up to a translation of  $\chi$ ) the following (invertible) map:

$$\begin{aligned} k_\ell &= (J_1^j - J_1^i) \chi_\ell, \ell \geq 2; \quad k_1 = \sum_{\ell=2}^n (J_\ell^i - J_\ell^j) \chi_\ell + \frac{1}{J_1^j - J_1^i} \left( \frac{\bar{\sigma}}{|\sigma|^2} w_0 + \sum_{\ell=1}^n J_\ell^i w_\ell \right); \\ \lambda_\ell &= \frac{(J_1^j - J_1^i)(J_\ell^i - J_\ell^j)}{\sigma_I J_\ell^i} (\sigma \chi_\ell)_I + w_\ell, \ell \geq 2; \quad \lambda_1 = \sum_{\ell=2}^n \left[ -\frac{(J_1^i - J_1^j)(J_\ell^i - J_\ell^j)}{\sigma_I J_1^i} (\sigma \chi_\ell)_I - \frac{J_\ell^i}{J_1^i} w_\ell \right]. \end{aligned} \quad (19)$$

To use (15) as before, we need the limit of  $T^{ij}$  for  $|\chi_p|$  large (and  $\chi_\ell$ ,  $\ell \neq p$ ,  $w_0$ ,  $w$  fixed); this turns out to depend on whether for some  $r \neq i, j$  we have

$$(J_1^r - J_1^j)(J_p^i - J_p^j) = (J_1^i - J_1^j)(J_p^r - J_p^j). \quad (20)$$

For brevity we consider only two cases (the only ones arising in the study of (2) - see the appendix): case I-relation (20) does not hold for any distinct  $i, j, r$  and any  $p \neq 1$ ; and case II-relation (20) holds for all  $i, j, r, p$  (in other words, the vectors  $J^1, \dots, J^m$  all lie on the same line in  $\mathbb{R}^n$ ). In the generic case I we have

$$\lim_{|\chi_p| \rightarrow \infty} T_\sigma^{ij}(\chi, w_0, w) = \hat{Q}^{ij}(w_0, w) \quad (21)$$

and (18)<sub>p1</sub><sup>ij</sup> becomes

$$I_p^{ij}[T_\sigma](\chi, w_0, w) \doteq T_\sigma^{ij}(\chi, w_0, w) - \frac{1}{\pi} \iint \frac{N_{p1}^{ij}[T_\sigma](\chi', w_0, w)}{\chi_p - \chi'_p} d\chi'_{R_p} d\chi'_{I_p} = \hat{Q}^{ij}(w_0, w), \quad (22)_I$$

where

$$\hat{Q}^{ij}(w_0, w) = \iint e^{-i(x_0 w_0 + x \cdot w)} Q^{ij}(x_0, x) dx_0 dx \quad \text{and } x' = (x_2, \dots, x_p', \dots, x_n).$$

If (20) holds for some  $r \neq j$  then (21) need not be true (see (7), (8), (11)).

In case II we have  $\frac{\partial T^{ij}}{\partial \bar{x}_p} \equiv 0$  for all  $p$ ,  $2 \leq p \leq n$ ; this, together with Liouville's

theorem, allows us to replace  $(22)_I$  by

$$T_\sigma^{ij}(\chi, w_0, w) = T_\sigma^{ij}(w_0, w). \quad (22)_{II}$$

In case I we conjecture (as in [2]) that the main condition needed to characterize the scattering data is that  $I_p^{ij}[T_\sigma](\chi, w_0, w)$  be independent of  $\chi$  and  $p$  and have suitable decay properties in  $(w_0, w)$ ; furthermore, given a  $T_\sigma$  which passes this admissibility test we can (re)construct a local potential  $Q$  simply as the inverse Fourier transform of  $I[T]$ .

From  $(22)_{II}$  we see that  $T^{ij}$  is completely determined by its values on the  $(n+1)$ -dimensional surface  $\chi = x_0$ ; the analogue of this in case I is the following: given  $T_\sigma^{ij}(x_0, w_0, w) = G^{ij}(w_0, w)$ ,  $1 \leq i, j \leq m$  we have (from  $(22)_I$ )

$$T_\sigma^{ij}(\chi, w_0, w) = G^{ij}(w_0, w) + \frac{1}{\pi} \iint \left[ \frac{N_{p1}^{ij}[T_\sigma](\chi', w_0, w)}{x_p - x_p'} - \frac{N_{p1}^{ij}[T_\sigma](x_0, w_0, w)}{x_{0p} - x_p'} \right] dx_{R_p}' dx_{I_p}' \quad (23)$$

which (at least for small  $G$ ) could be solved to find  $T$  everywhere.

### 3. The case $\sigma = -1$ .

If we formally substitute  $\sigma = -1$  in (6) we find that, away from the hyperplanes  $\Sigma_{ij} = \{k \in \mathbb{C}^n : \sum_{\ell=1}^n (J_\ell^i - J_\ell^j) k_{I_\ell} = 0\}$  the eigenfunction  $\mu_{-1}(x_0, x, k)$  is well-defined and holomorphic. Thus it appears that the inverse problem for the hyperbolic system  $L_{-1}$  could be regarded as a Riemann-Hilbert problem with data on the hyperplanes  $\Sigma_{ij}$ ,  $1 \leq i < j \leq m$ . We prefer to obtain an inversion procedure from our results for  $\sigma_I \neq 0$ . There seems to be little advantage in

studying the limit of  $\mu_\sigma(x_0, x, k)$  as  $\sigma \rightarrow -1$  (it leads us back to an analysis of singularities on the hyperplanes  $\Sigma_{ij}$ ); we work instead with the limit of  $\mu_\sigma(x_0, x, k_R, \sigma_I k_I)$ , with  $k_I$  now playing the role of a parameter. From (6) we find

$$\lim_{\sigma \rightarrow -1+i0} G_\sigma(x_0, x, k_R, \sigma_I k_I) = G_L(x_0, x, k_R, k_I)$$

$$= \frac{-i}{(2\pi)^{n+1}} \iint_{\mathbb{R}^{n+1}} \left\{ \frac{\theta(\sum_{\ell=1}^n [J_\ell^i \xi_\ell + (k_{R\ell} - k_{I\ell})(J_\ell^i - J_\ell^j)])}{\xi_0 - \sum_{\ell=1}^n [J_\ell^i \xi_\ell + k_{R\ell}(J_\ell^i - J_\ell^j)] + i0} + \frac{\theta(-\sum_{\ell=1}^n [J_\ell^i \xi_\ell + (k_{R\ell} - k_{I\ell})(J_\ell^i - J_\ell^j)])}{\xi_0 - \sum_{\ell=1}^n [J_\ell^i \xi_\ell + k_{R\ell}(J_\ell^i - J_\ell^j)] - i0} \right\} \times$$

$$\times e^{i(x_0 \xi_0 + x \cdot \xi)} d\xi_0 d\xi, \quad (24)$$

with  $\theta(\cdot)$  the Heaviside function; correspondingly,  $\lim_{\sigma \rightarrow -1+i0} \mu_\sigma(x_0, x, k_R, \sigma_I k_I) = \mu_L(x_0, x, k_R, k_I)$  where  $\mu_L$  solves the integral equation  $\mu_L = I + \tilde{G}_L(Q\mu_L)$ . From (24) we see that  $\mu_L(x_0, x, k_R, k_I)$  is a solution of

$$\frac{\partial \mu}{\partial x_0} - \sum_{\ell=1}^n J_\ell \frac{\partial \mu}{\partial x_\ell} - i \sum_{\ell=1}^n k_{R\ell} [J_\ell, \mu] = Q\mu \quad (25)$$

for every value of the parameter  $k_I$  in  $\mathbb{R}^n$ . Our scattering data is now

$$T_L^{ij}(k_R, k_I, \lambda) = \iint_{\mathbb{R}^{n+1}} e^{-i\beta_L^{ij}(x_0, x, k_R, k_I, \lambda)} (Q(x_0, x)\mu_L(x_0, x, k_R, k_I))^{ij} dx_0 dx \quad (26)$$

with  $\beta_L^{ij}(x_0, x, k_R, k_I, \lambda) = \sum_{\ell=1}^n [(J_\ell^i - J_\ell^j)(x_0 k_{I\ell} - \frac{x_\ell}{J_\ell^i}(k_{R\ell} - k_{I\ell})) + (x_\ell + J_\ell^i x_0)\lambda_\ell]$ . Taking limits in (14) we find the reconstruction equations for  $\mu$ :

$$\mu_L(x_0, x, k_R, k_I) = I + \frac{i}{(2\pi)^{n+1}} \sum_{i,j} (J_p^i - J_p^j) \iiint \left[ \frac{\theta(k_{I\ell} - k_{I\ell}')}{k_{R\ell} - k_{R\ell}' + i0} + \frac{\theta(k_{I\ell}' - k_{I\ell})}{k_{R\ell}' - k_{R\ell} - i0} \right] \delta(\sum_{\ell} J_\ell^i \lambda_\ell) \times$$

$$\times T_L^{ij}(k_R, k_I, \lambda) e^{i\beta_L^{ij}(x_0, x, k_R', k_I', \lambda)} \mu_L(x_0, x, k_R', k_I', \lambda) E_{ij} d\lambda dk_{R\ell}' dk_{I\ell}', \quad (27)$$

where now  $(\hat{k}_L^{ij}(k_R, k_I, \lambda))_{R_\ell} = \frac{J_\ell^j}{J_1^i} k_{R_\ell} + \frac{J_\ell^i - J_\ell^j}{J_1^i} k_{I_\ell} + \lambda_\ell$  and  $(\hat{k}_L^{ij})_{I_\ell} = k_{I_\ell}$ .

To write the characterization equations for  $T_L^{ij}$  we introduce new variables (suggested by the limit of (19))  $(x_R, x_I, w_0, w) \in \mathbb{R}^{3n-1}$  to parametrize the hyperplane  $\Sigma J_\ell^i \lambda_\ell = 0$  in  $\mathbb{R}^{3n}$  as follows:

$$\begin{aligned} k_{R_\ell} &= (J_1^j - J_1^i) x_{R_\ell}, \quad \ell \geq 2; \quad k_{R_1} = \sum_{\ell=2}^n (J_\ell^i - J_\ell^j) x_{R_\ell} + \frac{1}{J_1^i - J_1^j} (w_0 - \sum_{\ell=1}^n J_\ell^i w_\ell) \\ k_{I_\ell} &= (J_1^j - J_1^i) x_{I_\ell}, \quad \ell \geq 2; \quad k_{I_1} = \sum_{\ell=2}^n (J_\ell^i - J_\ell^j) x_{I_\ell} + \frac{1}{J_1^i - J_1^j} w_0 \end{aligned} \quad (28)$$

$$\lambda_\ell = \frac{(J_1^j - J_1^i)(J_\ell^i - J_\ell^j)}{J_1^i} (x_{R_\ell} - x_{I_\ell}) + w_\ell, \quad \ell \geq 2; \quad \lambda_1 = \sum_{\ell=2}^n \left[ \frac{(J_1^i - J_1^j)(J_\ell^i - J_\ell^j)}{J_1^i} (x_{R_\ell} - x_{I_\ell}) - \frac{J_\ell^i}{J_1^i} w \right]$$

Then under the assumptions of case I in section 2, the limit of the equations (22)<sub>I</sub> yields:

$$T_L^{ij}(x_R, x_I, w_0, w) = \hat{Q}^{ij}(w_0, w) + \frac{1}{\pi} \iint \left[ \frac{\theta(x_{I_p} - x_{I_p}')}{x_{R_p} - x_{R_p}' + i0} + \frac{\theta(x_{I_p}' - x_{I_p})}{x_{R_p} - x_{R_p}' - i0} \right] N_{p1}^{ij}[T_L](x', w_0, w) dx_{R_p}' dx_{I_p}', \quad (29)_I$$

with  $N_{p1}[T_L]$  given by a slight modification of (18). In case II we have

$$T_L^{ij}(x_R, x_I, w_0, w) = T_L^{ij}(w_0, w). \quad (29)_{II}$$

As in section 2, we can now use (29)<sub>I</sub> to characterize admissible  $T_L$ , (re)construct  $Q$ , as well as recover  $T_L$  from data given on  $x_R = \text{const.}$ ,  $x_I = \text{const.}$

It should be pointed out that once the family of Green's functions  $G_L$  has been chosen, all the above results can be derived without recourse to our limiting arguments ( $\nabla_{k_I} \mu_L$  can be expressed in terms of  $\mu_L$  using the appropriate symmetry property of  $G_L$  and the analytic behaviour of  $\mu_L$  for  $k_I$  large - needed to establish (27) - follows from (24); these analytic properties together with the

compatibility requirements  $\frac{\partial^2 \mu}{\partial k_{I_r} \partial k_{I_p}} = \frac{\partial^2 \mu}{\partial k_{I_p} \partial k_{I_r}}$  imply (29)).

#### 4. Relation between $T_L$ and the Scattering Operator ( $\sigma = -1$ )

To fix notation we sketch an elementary definition of the scattering operator associated with  $L_{-1}$ . When  $Q \equiv 0$ , given  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , the solution of the Cauchy problem  $L_{-1}u(x_0, x) = 0$ ,  $u(0, x) = f(x)$  is  $u^i(x_0, x) = f^i(x_1 + x_0 J_1^i, \dots, x_n + x_0 J_n^i)$ ,  $1 \leq i \leq m$ , which we write as  $u(x_0, x) = f(x + x_0 J)$ . When  $Q$  is, say, smooth and of compact support, given any (reasonable)  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  there is a unique  $u$  solution of  $L_{-1}u = 0$  with  $u(x_0, x) = f(x + x_0 J)$  for  $x_0 \ll 0$ ; furthermore there is a unique  $g$  such that  $u(x_0, x) = g(x + x_0 J)$  when  $x_0 \gg 0$ . We write  $g = Sf$ . On the Fourier transform side  $S$  can be written as

$$\hat{S}f(\xi) = f(\xi) + \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} S(\xi, k_R) \hat{f}(k_R) dk_R. \quad (30)$$

The question we would like to address is how to recover  $T_L$  (and hence  $Q$ ) given  $S(\xi, k_R)$ . To relate  $T_L$  and  $S(\xi, k_R)$  it turns out that we need to relate  $\mu_L$  and the eigenfunction  $\mu(x_0, x, k_R)$  corresponding to the "Volterra" Green's function

$$G^{ij}(x_0, x, k_R) = \theta(x_0) \exp[-i \sum_{\ell=1}^n (x_\ell + x_0 J_\ell^j) k_{R_\ell}] \prod_{\ell=1}^n \delta(x_\ell + x_0 J_\ell^i). \quad (31)$$

We start with the identity

$$\mu_L - \mu = (\tilde{G}_L - \tilde{G})(Q\mu_L) + \tilde{G}(Q(\mu_L - \mu)), \quad (32)$$

write  $G_L^{ij} - G^{ij}$  as a superposition of  $\exp(i\beta_L^{ij})$  and use a suitable symmetry property of  $G$ . The main result is the following linear equation for  $T_L$  given  $S$ :

$$\begin{aligned} T_L^{ij}(k_R, k_I, \lambda) &= S^{ij}(\hat{k}_R^{ij}(k_R, k_I, \lambda), k_R) - \frac{1}{(2\pi)^n} \sum_{i'} \int_{\mathbb{R}^n} \theta\left(\sum_{\ell=1}^n J_\ell^{i'} \lambda_\ell\right) \times \\ &\times S^{i'i'}(\hat{k}_R^{ij}(k_R, k_I, \lambda) \hat{k}_R^{i'j}(k_R, k_I, \lambda')) T_L^{i'j}(k_R, k_I, \lambda') d\lambda', \end{aligned} \quad (33)$$

where  $\hat{k}_R^{ij}(k_R, k_I, \lambda)$  stands for the real part of  $\hat{k}_L^{ij}$ .

### 5. Applications to Nonlinear Equations

The equations (2) are the compatibility conditions (cf. [ ]) for the Lax pair:

$$L_\sigma \psi = 0 \quad \text{and} \quad \frac{\partial \psi}{\partial t} + \sum_{\ell=1}^n B_\ell \frac{\partial \psi}{\partial x_\ell} = A \psi; \quad (34)$$

the matrices  $B_\ell$ ,  $1 \leq \ell \leq n$ , are constant real diagonal and  $A^{ij}(t, x_0, x) = \frac{1}{\sigma} a_{ij} Q^{ij}(t, x_0, x)$  with  $a_{ij}$  given by (3). The restrictions imposed by (3) on the matrices  $J_\ell$ ,  $1 \leq \ell \leq n$ , are discussed in the appendix. To find the time dependence of the scattering data corresponding to (2), we set  $\psi = \mu \exp[i \sum_{\ell=1}^n k_\ell (x_\ell - \sigma x_0 J_\ell - t B_\ell)]$ ; then  $\mu$  satisfies (4) as well as

$$A\mu \doteq \frac{\partial \mu}{\partial t} + \sum_{\ell=1}^n B_\ell \frac{\partial \mu}{\partial x_\ell} + i \sum_{\ell=1}^n k_\ell [B_\ell, \mu] - A\mu = 0. \quad (35)$$

Applying the operator  $A$  to both sides of the equation (14) we find (when  $\sigma_I \neq 0$ )

$$-\frac{\partial T_\sigma^{ij}}{\partial t}(t, k, \lambda) = i \sum_{\ell=1}^n [B_\ell^j k_\ell - B_\ell^i \hat{k}_\ell^{ij}(k, \lambda)] T_\sigma^{ij}(t, k, \lambda). \quad (36)$$

For the case  $\sigma = -1$  the equations (obtained as limits of (36) or by a parallel derivation) are

$$\frac{\partial T_L^{ij}}{\partial t}(t, k_R, k_I, \lambda) = i \sum_{\ell=1}^n [B_\ell^j k_{R_\ell} - B_\ell^i \hat{k}_{R_\ell}^{ij}(k_R, k_I, \lambda)] T_L^{ij}(t, k_R, k_I, \lambda). \quad (37)$$

Thus, when  $\sigma = -1$ , we can apply the inverse scattering procedure together with (37) to construct the solution to the Cauchy problem for (2). Note that  $T_L(t, k_R, k_I, \lambda)$  as given by (37) satisfies the characterization equations if  $T_L(0, k_R, k_I, \lambda)$  does.

When  $\sigma_I \neq 0$  the Cauchy problem for (2) is ill-posed; however (by analogy to the corresponding linear problem) we can use inverse scattering to solve (2) as follows: given  $Q(0, x_0, x)$  it can be decomposed into  $Q_+(0, x_0, x) + Q_-(0, x_0, x)$

where  $Q_+(0, x_0, x)$  extends to a function  $Q_+(t, x_0, x)$  satisfying (2) in the half-space  $t > 0$ , while  $Q_-(0, x_0, x)$  extends to a function satisfying (2) in the half-space  $t < 0$ . Assume for simplicity that  $\sigma_I a_{ij} > 0$  for all  $i \neq j$ . Given  $Q$  define  $Q_+$  by  $\hat{Q}_+(0, w_0, w) = \theta(\bar{w}_0) \hat{Q}(0, w_0, w)$ . If  $T_+$  is the scattering transform of  $Q_+$  then from the direct problem we find  $T_+^{ij}(0, \chi, w_0, w) = 0$  for  $w_0 > 0$ ; thus for  $t > 0$  we can define (see (36))  $T_+^{ij}(t, \chi, w_0, w)$  by

$$\begin{aligned} T_+^{ij}(t, \chi, w_0, w) &= \exp[it \sum_{\ell=1}^n (B_{\ell}^j k_{\ell} - B_{\ell}^i \hat{k}_{\ell}^{ij})] T_+^{ij}(0, \chi, w_0, w) = (\text{see (3), (13) and (19)}) \\ &= \exp[it(\frac{a_{ij}}{\sigma} w_0 + \sum_{\ell=1}^n (a_{ij} j_{\ell}^i - B_{\ell}^i) w_{\ell})] T_+^{ij}(0, \chi, w_0, w). \end{aligned} \quad (38)$$

Since the expression in the exponential does not depend on  $\chi$  and since its real part is nonpositive if  $t > 0$ ,  $T_+^{ij}(t, \chi, w_0, w)$  satisfies the characterization equations (29) so inverse scattering should yield the desired potential  $Q_+(t, x_0, x)$ ; similarly we can construct  $Q_-(t, x_0, x)$  solution of (2) for  $t < 0$ .

#### Acknowledgements

It is a pleasure to thank A.S. Fokas for helpful conversations. This work was supported in part by the National Science Foundation under grant number DMS-8202117, the Office of Naval Research under grant number N00014-76-C-0867, the Air Force Office of Scientific Research under grant number AFOSR-84-0005 and by a University of Rochester faculty research grant.



# Appendix

We need to find the restrictions imposed on the choice of matrices  $J_\ell$ ,  $1 \leq \ell \leq n$ , by the existence of  $(a_{ij})$  and  $B_\ell$ ,  $1 \leq \ell \leq n$  satisfying (3).

If (3) holds then the matrix  $(a_{ij})$  is symmetric and

$$a_{ip} - a_{pj} = (a_{ij} - a_{pj}) \frac{J_\ell^j - J_\ell^i}{J_\ell^p - J_\ell^i} \text{ for every } \ell \text{ and every } i, j, p \text{ distinct.} \quad (A1)$$

(Conversely, if (A1) holds with  $(a_{ij})$  symmetric then  $B_\ell$ ,  $1 \leq \ell \leq n$  can be found so that (3) is satisfied.) Note that if  $a_{ip} \neq a_{pj}$ , (A1) forces  $J^i$ ,  $J^j$ ,  $J^p$  to be colinear. There are two cases:

I  $a_{ip} = a_{pj}$  for all  $i, j, p$  distinct. Then (A1) puts no restriction on  $J_\ell$ ; in particular they can be chosen so that (20) does not hold for any distinct  $i, j, r$  and  $p \neq 1$ . The system (2) is linear in this case.

II For some  $i_0, j_0, p_0$  distinct  $a_{i_0 p_0} \neq a_{p_0 j_0}$ . We show that in this case the vectors  $J^1, \dots, J^m$  must all be colinear. From (A1) we already know that  $J^{i_0}, J^{j_0}, J^{p_0}$  are colinear. For any  $r \neq i_0, j_0, p_0$  one of the following must be true

$$(i) a_{i_0 r} \neq a_{r j_0}, \quad (ii) a_{r i_0} \neq a_{i_0 p_0}, \quad (iii) a_{r j_0} \neq a_{j_0 p_0} \quad (A2)$$

(for if not  $a_{i_0 p_0} = a_{r i_0} = a_{r j_0} = a_{p_0 j_0}$  contradicting our assumption). In either of the possibilities (A2)  $J^r$  will be on the line passing through  $J^{i_0}, J^{p_0}, J^{j_0}$ ; this will be true for any  $r$ ,  $1 \leq r \leq m$ . (Conversely, given  $J^1, J^2, \dots, J^m$  colinear with  $J_\ell^i \neq J_\ell^j$  we can construct  $(a_{ij})$  symmetric satisfying II and (A1)).

It follows that whenever (2) is not linear, the matrix having  $J^1, J^2, \dots, J^m$  as rows has rank at most 2; thus if  $n \geq 3$  its columns (the diagonals of the matrices  $J_\ell$  in (1)) must be linearly dependent and then the inverse scattering problem for  $L_\sigma$  can also be solved by reducing it to a lower dimensional one. On the other hand, since the characterization equations are trivial (i.e.  $N(T) = 0$ ) in this case, it seems reasonable to expect that other (possibly non-local) nonlinear equations

can be found which would be compatible with (22)<sub>II</sub>.

#### References

- [1] A.S. Fokas and M.J. Ablowitz, Institute for Nonlinear Studies Report #24 (to appear in J. Math. Phys.); A.S. Fokas, Phys. Rev. Lett. 51, 3 (1983); A.S. Fokas and M.J. Ablowitz, Phys. Rev. Lett. 51, 7 (1983); see also the review articles by M.J. Ablowitz and A.S. Fokas in "Nonlinear Phenomena", Proceedings, Oaxtepec, Mexico 1982, pp. 3-24 and 137-183, Edited by K.B. Wolf, Lecture Notes in Physics #189, Springer. For earlier work on these subjects see D.J. Kaup, Physica 1D, 45 (1980) and L.P. Niznik and V.G. Tarasov, Soviet Math. Dokl. Vol. 18 No. 2, 397 (1977).
- [2] A.I. Nachman and M.J. Ablowitz, Institute for Nonlinear Studies Report, #37 (to appear in Stud. Appl. Math.).
- [3] M.J. Ablowitz and R. Haberman, Phys. Rev. Lett. 35, 1185 (1975); B.G. Konopelchenko, Phys. Lett. 93A, 442 (1983).

**END**

**FILMED**

**1-85**

**DTIC**